

[Go to Product page](#)

Datasheet for ABIN745253

**anti-RPS6KA2 antibody (pSer360, pThr356)**

## Overview

Quantity:	100 µL
Target:	RPS6KA2
Binding Specificity:	pSer360, pThr356
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RPS6KA2 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human RSK3 around the phosphorylation site of Thr356/Ser360
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

## Target Details

Target:	RPS6KA2
Alternative Name:	RSK3 + ( <a href="#">RPS6KA2 Products</a> )

## Target Details

Background: Synonyms: 90 kDa ribosomal protein S6 kinase 2, HU 2, KS6A2\_HUMAN, MAP kinase activated protein kinase 1c, MAP kinase-activated protein kinase 1c, MAPK-activated protein kinase 1c, MAPKAP kinase 1c, MAPKAPK-1c, MAPKAPK1C, p90-RSK 2, p90RSK 2, p90RSK2, pp90RSK 3, pp90RSK3, Ribosomal protein S6 kinase alpha-2, Ribosomal S6 kinase 3, RPS6KA2, RSK 3, RSK-3, S6K alpha 2, S6K alpha, S6K-alpha-2.

Background: RSK3(Ribosomal S6 kinase 3)is a member of a family of 90- kDa ribosomal protein S6 kinases, which includes Rsk1, Rsk2 and Rsk3. These are broadly expressed serine/threonine protein kinases that are activated in response to mitogenic stimuli, including extracellular signal-regulated protein kinases Erk1 and Erk2. Rsk3 is a distinct isoform of p90Rsk that translocates to the cell nucleus, phosphorylates potential nuclear targets and may have a unique upstream activator. Several sites, such as Ser380, Thr359, Ser363 and Thr573, are important for its activation. Active Rsks appear to play a major role in transcriptional regulation by translocating to the nucleus and phosphorylating c Fos and CREB.

Gene ID: 6196

Pathways: [MAPK Signaling](#), [Neurotrophin Signaling Pathway](#), [Regulation of Systemic Arterial Blood Pressure by Hormones](#), [Activation of Innate immune Response](#), [Toll-Like Receptors Cascades](#)

## Application Details

Application Notes: ELISA 1:500-1000  
IHC-P 1:200-400  
IHC-F 1:100-500  
IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

## Handling

---

handled by trained staff only.

---

Storage: 4 °C,-20 °C

---

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

---

Expiry Date: 12 months