

Datasheet for ABIN7453107
anti-IDH3B antibody (AA 25-75)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	IDH3B
Binding Specificity:	AA 25-75
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)

Product Details

Purpose:	Rabbit anti-IDH3B Antibody, Affinity Purified
Immunogen:	Between AA 25 and 75
Isotype:	IgG
Predicted Reactivity:	Pig,Orangutan
Purification:	Affinity Purified

Target Details

Target:	IDH3B
Alternative Name:	IDH3B (IDH3B Products)
Background:	Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+)

Target Details

as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. Isocitric dehydrogenase subunit beta (IDH3B) is the beta subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase [taken from NCBI Entrez Gene (Gene ID: 3420)].

Gene ID: 3420

UniProt: [O43837](#)

Application Details

Application Notes: IP: Not recommended
WB: 1:2,000 - 1:10,000

Restrictions: For Research Use only

Handling

Concentration: 1000 µg/mL

Buffer: Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Expiry Date: 12 months