antibodies .- online.com





anti-Cyclin T1 antibody (AA 625-675)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	20 μg
Target:	Cyclin T1 (CCNT1)
Binding Specificity:	AA 625-675
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin T1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	Rabbit anti-Cyclin T1 Antibody, Affinity Purified
Immunogen:	between AA 625 and 675
Isotype:	IgG
Predicted Reactivity:	Chimpanzee
Purification:	Affinity Purified

Target Details

Target:	Cyclin T1 (CCNT1)
Alternative Name:	Cyclin T1 (CCNT1 Products)

Target Details

Target Details		
Background:	Background: Cyclin T1 is a regulatory subunit of the cyclin-dependent kinase pair (CDK9/cyclin-T1) complex, also called positive transcription elongation factor B (P-TEFb), which is proposed to facilitate the transition from abortive to productive elongation by phosphorylating the CTD (carboxy-terminal domain) of the large subunit of RNA polymerase II (RNA Pol II). In case of HIV or SIV infections, Cyclin T1 binds to the transactivation domain of the viral nuclear transcriptional activator, Tat, thereby increasing Tat's affinity for the transactivating response RNA element (TAR RNA) [taken from the Universal Protein Resource (UniProt) 060563].	
Gene ID:	904	
NCBI Accession:	NP_001231	
UniProt:	O60563	
Application Details		
Application Notes:	IHC: 1:100 - 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. IP: 2 - 10 µg/mg lysate WB: 1:2,000 - 1:10,000	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Expiry Date:	12 months	