

## Datasheet for ABIN7453646 anti-STAT2 antibody (AA 725-775)



## Go to Product page

( )	ver	~\ /I	~ 1 A
	$\vee \square$	\/ I	ᆸᇄ
$\sim$	v Cı	V I	~ v ·

Background:

Quantity:	20 μg
Target:	STAT2
Binding Specificity:	AA 725-775
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed
	Paraffin-embedded Sections) (IHC (fp))
Product Details	
Purpose:	Rabbit anti-STAT2 Antibody, Affinity Purified
Immunogen:	between AA 725 and 775
Isotype:	IgG
Purification:	Affinity Purified
Target Details	
Target:	STAT2
Alternative Name:	STAT2 (STAT2 Products)

Background: Signal transducer and activator of transcription 2 (STAT2) is a signal transducer

Expiry Date:

12 months

Target Details	
	and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state [taken from the Universal Protein Resource (UniProt) P52630].
Gene ID:	6773
NCBI Accession:	NP_005410
UniProt:	P52630
Pathways:	JAK-STAT Signaling, Hepatitis C, CXCR4-mediated Signaling Events
Application Details	
Application Notes:	IHC: 1:100 - 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue
	sections.  IP: 2 - 10 µg/mg lysate  WB: 1:2,000 - 1:10,000
Restrictions:	IP: 2 - 10 μg/mg lysate
Restrictions: Handling	IP: 2 - 10 μg/mg lysate WB: 1:2,000 - 1:10,000
	IP: 2 - 10 μg/mg lysate WB: 1:2,000 - 1:10,000
Handling	IP: 2 - 10 µg/mg lysate  WB: 1:2,000 - 1:10,000  For Research Use only
Handling Concentration:	IP: 2 - 10 μg/mg lysate WB: 1:2,000 - 1:10,000  For Research Use only  200 μg/mL
Handling Concentration: Buffer:	IP: 2 - 10 μg/mg lysate  WB: 1:2,000 - 1:10,000  For Research Use only  200 μg/mL  Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide