

Datasheet for ABIN7453647 anti-STAT2 antibody (AA 801-851)



Go to Product page

\sim				
()\	ve	r\/		Λ/
\cup	$V \subset$	ı vı	\Box	٧V

Quantity: 20 µg Target: STAT2 Binding Specificity: AA 801-851 Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purfication: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products) Background: Background: Signal transducer and activator of transcription 2 (STAT2) is a signal transducer	Overview		
Binding Specificity: AA 801-851 Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 Products)	Quantity:	20 μg	
Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Target:	STAT2	
Host: Rabbit Clonality: Polyclonal Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Binding Specificity:	AA 801-851	
Clonality: Polyclonal Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Reactivity:	Human	
Conjugate: This STAT2 antibody is un-conjugated Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Host:	Rabbit	
Application: Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Clonality:	Polyclonal	
Paraffin-embedded Sections) (IHC (fp)) Product Details Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Conjugate:	This STAT2 antibody is un-conjugated	
Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed	
Purpose: Rabbit anti-STAT2 Antibody, Affinity Purified Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)		Paraffin-embedded Sections) (IHC (fp))	
Immunogen: between AA 801 and 851 Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Product Details		
Isotype: IgG Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Purpose:	Rabbit anti-STAT2 Antibody, Affinity Purified	
Purification: Affinity Purified Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Immunogen:	between AA 801 and 851	
Target Details Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Isotype:	IgG	
Target: STAT2 Alternative Name: STAT2 (STAT2 Products)	Purification:	Affinity Purified	
Alternative Name: STAT2 (STAT2 Products)	Target Details		
	Target:	STAT2	
Background: Background: Signal transducer and activator of transcription 2 (STAT2) is a signal transduc	Alternative Name:	STAT2 (STAT2 Products)	
	Background:	Background: Signal transducer and activator of transcription 2 (STAT2) is a signal transducer	

rarget Details		
	and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta).	
	Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are	
	activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs	
	dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor,	
	that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate	
	the transcription of interferon stimulated genes, which drive the cell in an antiviral state [taken	
	from the Universal Protein Resource (UniProt) P52630].	
Gene ID:	6773	
NCBI Accession:	NP_005410	
UniProt:	P52630	
Pathways:	JAK-STAT Signaling, Hepatitis C, CXCR4-mediated Signaling Events	
Application Details		
Application Notes:	IHC: 1:100 - 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue	
	sections.	
	IP: 2 - 10 μg/mg lysate	
	WB: 1:2,000 - 1:10,000	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	

4°C

12 months

Storage:

Expiry Date: