.-online.com antibodies

Datasheet for ABIN7453800 anti-TDP1 antibody (AA 1-50)



Overview

Quantity:	20 µg
Target:	TDP1
Binding Specificity:	AA 1-50
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TDP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	Rabbit anti-TDP1 Antibody, Affinity Purified
Immunogen:	between AA 1 and 50
Isotype:	IgG
Purification:	Affinity Purified

Target Details

Target:	TDP1
Alternative Name:	TDP1 (TDP1 Products)
Background:	Background: Tyrosyl-DNA phosphodiesterase 1 (TDP1) is an enzyme that participates in the

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7453800 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	repair of DNA strand breaks associated with topoisomerase 1 (Top1) complexes. TDP1
	catalyzes the removal of Top1 from DNA via the hydrolysis of the phosphodiester bond
	between the tyrosine residue of Top1 and the 3' phosphate of DNA. Removal of Top1 is then
	followed by DNA repair. TDP1 has been proposed to play more general roles in DNA damage
	repair independent of the Top1 complex. TDP1 has been shown to associate with DNA ligase
	IIIalpha, a component of the DNA single-stranded break repair (SSBR) machinery. It is also able
	to remove glycolate from 3'-phosphoglycolate termini of double-stranded breaks and a variety
	of other 3'- adducts. Defects in TDP1 are associated with spinocerebellar ataxia autosomal
	recessive with axonal neuropathy (SCAN1), a disease associated with peripheral axonal motor
	and sensory neuropathy and distal muscular atrophy.
Gene ID:	55775
NCBI Accession:	NP_001008744
UniProt:	Q9NUW8
Application Details	
Application Notes:	IHC: 1:100 - 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue
	sections.
	IP: 2 - 5 µg/mg lysate
	WB: 1:2,000 - 1:10,000
Restrictions:	For Research Use only
Handling	
Concentration:	200 µg/mL
Buffer:	Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months