



[Go to Product page](#)

Datasheet for ABIN7454057
anti-Apo-B100 antibody (FITC)

Overview

Quantity:	0.5 mg
Target:	Apo-B100
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This Apo-B100 antibody is conjugated to FITC

Product Details

Purpose:	Goat anti-Human Apo B-100 [F06] Antibody FITC Conjugated
Immunogen:	Human Apolipoprotein B100
Clone:	F06
Isotype:	IgG

Target Details

Target:	Apo-B100
Alternative Name:	ApoB-100 (Apo-B100 Products)
Background:	Background: ApoB exists in human plasma in two isoforms, ApoB-48 (Chen et al., 1987) and Apo B-100 (Wei et al., 1985, Yang et al., 1986a, 1989a,b, 1990, Chen et al., 1986, Yang et al., 1990, Yang and Pownall, 1992). Apo B-100 is the major physiological ligand for the LDL receptor. Apo B-100 is a large monomeric protein, containing 4536 amino acids (m.w. 515 kDa, Yang et al., 1986b). Apo B-100 is synthesized in the liver and is required for the assembly of

Target Details

VLDL. It is found in LDL and VLDL after the removal of the Apo A, E and C. Apo B-48 is present in chylomicrons and their remnants. It is essential for the intestinal absorption of dietary lipids. Apo B levels correlate with the risk of coronary disease. The Apo B protein is directly involved in the retention of LDL with the arterial wall (Olofsson and Boren, 2012). Apo B-48 is synthesized in the small intestine. It comprises approximately half of the N-terminal region of ApoB-100 and is the result of posttranscriptional mRNA editing by a stop codon in the intestine not found in the liver.

Gene ID: 338

NCBI Accession: [NP_000375](#)

UniProt: [P04114](#)

Application Details

Restrictions: For Research Use only

Handling

Concentration: 1 mg/mL

Buffer: 75 mM Sodium Phosphate, 75 mM NaCl, 0.5 mM EDTA, 0.02 % Sodium azide, pH 7.2

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Expiry Date: 12 months