

Datasheet for ABIN7454118

FGL2 Protein (AA 197-432) (His-Avi-DYKDDDDK Tag, Biotin)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μg
Target:	FGL2
Protein Characteristics:	AA 197-432
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FGL2 protein is labelled with His-Avi-DYKDDDDK Tag,Biotin.

Product Details

Purpose:	Biotinylated Mouse FGL2 Protein	
Sequence:	Pro197-Pro432	
Characteristics:	Recombinant Biotinylated Mouse FGL2 Protein is expressed from HEK293 with His tag and Avi tag and Flag tag at the N-Terminus.It contains Pro197-Pro432.	
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC	
Sterility:	0.22 µm filtered	
Endotoxin Level:	Less than 1EU per µg by the LAL method.	

Target Details

Target:	FGL2	
Alternative Name:	FGL2 (FGL2 Products)	

Target Details

Expiry Date:

12 months

important regulatory functions in both innate and adaptive immune responses. FGL2 is overexpressed in glioma, and its expression level is negatively associated with the progno glioma patients. Molecular Weight: 31.48 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result. UniProt: P12804 Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after	rai got Botano	
result. UniProt: P12804 Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Background:	overexpressed in glioma, and its expression level is negatively associated with the prognosis of
Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Molecular Weight:	31.48 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.
Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	UniProt:	P12804
Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Application Details	
Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Restrictions:	For Research Use only
Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/r recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Handling	
recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Format:	Lyophilized
protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein in	Storage:	-20 °C,-80 °C
	Storage Comment:	reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into