Datasheet for ABIN7454202

antibodies.com

PDGFRB Protein (AA 32-532) (His tag)



Overview

Quantity:	100 µg
Target:	PDGFRB
Protein Characteristics:	AA 32-532
Origin:	Dog
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PDGFRB protein is labelled with His tag.

Product Details

Purpose:	Canine PDGF R beta/CD140b Protein
Sequence:	Gly32-Val532
Characteristics:	Recombinant Canine PDGF R beta/CD140b Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Gly32-Val532.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

Target Details

Target:	PDGFRB
Alternative Name:	PDGF R beta (PDGFRB Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7454202 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	PDGF is a major serum mitogen that can exist as a homo or hetero-dimeric protein consisting
	of disulfide-linked PDGF-A and PDGF-B chains. The PDGF-AA,PDGF-BB and PDGF-AB isoforms
	have been shown to bind to two distinct cell surface PDGF receptors with different affinities.
	Where as PDGF R alpha binds all three PDGF isoforms with high affinity, PDGF R beta binds
	PDGF-BB only with high-affinity. PDGF R plays an essential role in blood vessel development by
	promoting proliferation, migration and recruitment of pericytes and smooth muscle cells to
	endothelial cells. Plays a role in the migration of vascular smooth muscle cells and the
	formation of neointima at vascular injury sites.
Molecular Weight:	57.25 kDa. Due to glycosylation, the protein migrates to 70-100 kDa based on Tris-Bis PAGE
	result.
UniProt:	Q6QNF3
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling
	Pathway, Inositol Metabolic Process, Glycosaminoglycan Metabolic Process, Smooth Muscle
	Cell Migration, Platelet-derived growth Factor Receptor Signaling

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7454202 | 01/18/2024 | Copyright antibodies-online. All rights reserved.