antibodies.com

Datasheet for ABIN7454253 FGF2 Protein (AA 135-288)



Overview

| Quantity: | 100 µg |
|--------------------------|----------------------------|
| Target: | FGF2 |
| Protein Characteristics: | AA 135-288 |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |

Product Details

| Purpose: | Human FGF basic (154aa) Protein |
|------------------------------|---|
| Sequence: | Ala135-Ser288 |
| Characteristics: | Recombinant Human FGF basic (154aa) Protein is expressed from E.coli without tag.lt contains Ala135-Ser288. |
| Purity: | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per μ g by the LAL method. |
| Biological Activity Comment: | Immobilized Human FGF basic (154aa) at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Human Glypican 1, His Tag with the EC50 of 5.2ng/ml determined by ELISA. See testing image for detail. |

Target Details

Target:

FGF2

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7454253 | 01/18/2024 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|---------------------|--|
| Alternative Name: | FGF basic (FGF2 Products) |
| Background: | FGF basic is a member of the FGF family of at least 23 related mitogenic proteins which show 35-60 % amino acid conservation. FGF acidic and basic, unlike the other members of the family, lack signal peptides and are apparently secreted by mechanisms other than the classical protein secretion pathway. |
| Molecular Weight: | 17.12 kDa same as Tris-Bis PAGE result. |
| Pathways: | RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, C21-Steroid Hormone Metabolic Process, Inositol Metabolic Process, Glycosaminoglycan Metabolic Process, Protein targeting to Nucleus, S100 Proteins |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water. |
| Buffer: | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Expiry Date: | 12 months |