

[Go to Product page](#)

Datasheet for ABIN7454453 **anti-EAF2 antibody (AA 150-200)**

Overview

| | |
|----------------------|---|
| Quantity: | 100 µg |
| Target: | EAF2 |
| Binding Specificity: | AA 150-200 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EAF2 antibody is un-conjugated |
| Application: | Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)) |

Product Details

| | |
|---------------|--|
| Purpose: | Rabbit anti-EAF2 Antibody, Affinity Purified |
| Immunogen: | between AA 150 and 200 |
| Isotype: | IgG |
| Purification: | Affinity Purified |

Target Details

| | |
|-------------------|--|
| Target: | EAF2 |
| Alternative Name: | EAF2 (EAF2 Products) |
| Background: | Background: ELL-associated factor 2 (EAF2) acts as a transcriptional transactivator of TCEA1 |

Target Details

elongation activity. It is also a transcriptional transactivator of ELL (eleven nineteen lysine-rich leukemia) 1 and ELL2 elongation activities. EAF2 has been shown to be a potent inducer of apoptosis in prostatic and non-prostatic cell lines and to inhibit prostate tumor growth in vivo [taken from the Universal Protein Resource (UniProt) Q96CJ1].

Gene ID: 55840

NCBI Accession: [NP_060926](#)

UniProt: [Q96CJ1](#)

Application Details

Application Notes: IHC: 1:500 - 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.

IP: 2 - 5 µg/mg lysate

WB: Not recommended

Restrictions: For Research Use only

Handling

Concentration: 1000 µg/mL

Buffer: Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Expiry Date: 12 months