antibodies -online.com





Datasheet for ABIN7454828 anti-MED13L antibody (AA 550-600)

Go to Product page

/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	Λ
\sim	' V C	1 V		v v

Quantity:	20 μg
Target:	MED13L
Binding Specificity:	AA 550-600
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED13L antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	Rabbit anti-MED13L Antibody, Affinity Purified	
Immunogen:	between AA 550 and 600	
Isotype:	IgG	
Purification:	Affinity Purified	

Target Details

Target:	MED13L
Alternative Name:	MED13L (MED13L Products)
Background: Background: Mediator complex subunit 13-like is a component of the Mediator complex	

coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent		
genes. Mediator functions as a bridge to convey information from gene-specific regulatory		
proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to		
promoters by direct interactions with regulatory proteins and serves as a scaffold for the		
assembly of a functional preinitiation complex with RNA polymerase II and the general		
transcription factors. This subunit may specifically regulate transcription of targets of the Wnt		
signaling pathway and SHH signaling pathway. [taken from the Universal Protein Resource		
(UniProt) Q71F56].		

Gene ID:	23389
NCBI Accession:	NP_056150
UniProt:	Q71F56

Pathways: Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes:	IHC: 1:200 to 1:1000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE	
	tissue sections.	
	IP: 2 - 10 μg/mg lysate	
	WB: 1:2,000 - 1:10,000	

Restrictions: For Research Use only

Handling

Concentration:	200 μg/mL
Buffer:	Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months