

Datasheet for ABIN7454851 anti-EYA2 antibody (AA 100-150)



Go to Product page

Overview	
Quantity:	100 μg
Target:	EYA2
Binding Specificity:	AA 100-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EYA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))
Product Details	
Purpose:	Rabbit anti-EYA2 Antibody, Affinity Purified
Immunogen:	between AA 100 and 150

Purpose:	Rabbit anti-EYA2 Antibody, Affinity Purified
Immunogen:	between AA 100 and 150
Isotype:	IgG
Predicted Reactivity:	Bovine
Purification:	Affinity Purified

Target Details

Target:	EYA2
Alternative Name:	EYA2 (EYA2 Products)

Target Details

rarget Details	
Background:	Background: EYA2 (eyes absent homolog 2) functions as a tyrosine phosphatase that specifically dephosphorylates 'Tyr-142' of histone H2AX. EYA2 promotes efficient DNA repair by dephosphorylating H2AX and promoting the recruitment of DNA repair complexes containing MDC1. EYA2 coactivates SIX1. Together with SIX1 and DACH2 EYA2 seems to be involved in myogenesis. EYA2 may be involved in development of the eye [taken from the Universal Protein Resource (UniProt) 000167].
Gene ID:	2139
NCBI Accession:	NP_005235
UniProt:	000167
Application Details	
Application Notes:	IHC: 1:500 - 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. IP: 2 - 5 µg/mg lysate WB: 1:2,000 - 1:10,000
Restrictions:	For Research Use only
Handling	
Concentration:	1000 μg/mL
Buffer:	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months