

Datasheet for ABIN7454967
anti-PSMD7 antibody (AA 150-200)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PSMD7
Binding Specificity:	AA 150-200
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMD7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	Rabbit anti-PSMD7 Antibody, Affinity Purified
Immunogen:	Between AA 150 and 200
Isotype:	IgG
Predicted Reactivity:	D. melanogaster,Bovine
Purification:	Affinity Purified

Target Details

Target:	PSMD7
Alternative Name:	PSMD7 (PSMD7 Products)

Target Details

Background:	Background: Proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (PSMD7) is a subunit of the 26S proteasome, a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMD7 is a non-ATPase subunit of the 19S regulator. [taken from NCBI Entrez Gene (Gene ID: 5713)].
Gene ID:	5713
NCBI Accession:	NP_002802
UniProt:	P51665
Pathways:	Mitotic G1-G1/S Phases , DNA Replication , Synthesis of DNA , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	IHC: 1:500 - 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. IP: 2 - 10 µg/mg lysate WB: 1:2,000 - 1:10,000
Restrictions:	For Research Use only

Handling

Concentration:	1000 µg/mL
Buffer:	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months