

Datasheet for ABIN7454976
anti-PSMD2 antibody (AA 50-100)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PSMD2
Binding Specificity:	AA 50-100
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

Product Details

Purpose:	Rabbit anti-PSMD2 Antibody, Affinity Purified
Immunogen:	Between AA 50 and 100
Isotype:	IgG
Predicted Reactivity:	Rat,Bovine,Orangutan
Purification:	Affinity Purified

Target Details

Target:	PSMD2
Alternative Name:	PSMD2 (PSMD2 Products)
Background:	Background: PSMD2 [proteasome (prosome, macropain) 26S subunit, non-ATPase, 2] is a

Target Details

subunit of the 26 S proteasome. The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMD2 is a subunit of the 19S regulator lid, which is responsible for substrate recognition and binding [taken from NCBI Entrez Gene (Gene ID: 5708)].

Gene ID:	5708
NCBI Accession:	NP_002799
UniProt:	Q13200
Pathways:	Mitotic G1-G1/S Phases , DNA Replication , Synthesis of DNA , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	IHC: 1:500 - 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. IP: 2 - 10 µg/mg lysate WB: 1:1,000 - 1:5,000
Restrictions:	For Research Use only

Handling

Concentration:	1000 µg/mL
Buffer:	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months