

#### Datasheet for ABIN7455140

# Recombinant anti-MED1 antibody (AA 1550-1581)



#### Overview

Quantity:	100 μL	
Target:	MED1	
Binding Specificity:	AA 1550-1581	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Antibody Type:	Recombinant Antibody	
Clonality:	Monoclonal	
Conjugate:	This MED1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp)), Flow Cytometry (FACS)	

#### **Product Details**

Purpose:	Rabbit anti-MED1 Recombinant Monoclonal Antibody [BLR037F]	
Immunogen:	between AA 1550 and 1581 (C-term)	
Clone:	BLR037F	
Isotype:	IgG	
Predicted Reactivity:	Orangutan	

### **Target Details**

Target:	MED1	
Alternative Name:	MED1 (MED1 Products)	
Background:	Background: MED1 is a component of the Mediator complex, a coactivator involved in the	
	regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as	
	a bridge to convey information from gene-specific regulatory proteins to the basal RNA	
	polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions	
	with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation	
	complex with RNA polymerase II and the general transcription factors [taken from the Universal	
	Protein Resource (UniProt) Q15648].	
Gene ID:	5469	
NCBI Accession:	NP_004765	
UniProt:	Q15648	
Pathways:	Nuclear Receptor Transcription Pathway, Intracellular Steroid Hormone Receptor Signaling	
	Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Nuclear Hormone	
	Receptor Binding, Chromatin Binding, Regulation of Lipid Metabolism by PPARalpha	
Application Details		
Application Notes:	IHC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue	
	sections.	
	IP: 20 µL/1 mg lysate	
	IP: 20 $\mu$ L/1 mg lysate ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue	
	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue	
Restrictions:	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.	
	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  WB: 17:40:00	
Restrictions: Handling Concentration:	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  WB: 17:40:00  For Research Use only	
Handling Concentration:	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  WB: 17:40:00  For Research Use only  100 μg/mL	
Handling Concentration: Buffer:	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  WB: 17:40:00  For Research Use only  100 µg/mL  Borate Buffered Saline (BBS) pH 8.2 with 0.1 % BSA and 0.09 % Sodium Azide	
Handling	ICC: 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.  WB: 17:40:00  For Research Use only  100 μg/mL	

## Handling

Storage:	4 °C	
Expiry Date:	12 months	