

Datasheet for ABIN7456501

anti-CD161 antibody (APC)



Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

0.10.11011			
Quantity:	20 tests		
Target:	CD161 (KLRB1)		
Reactivity:	Human		
Host:	Mouse		
Clonality:	Monoclonal		
Conjugate:	This CD161 antibody is conjugated to APC		
Application:	Flow Cytometry (FACS)		
Product Details			
Clone:	HP-3G10		
Isotype:	IgG1, kappa		
Target Details			
Target:	CD161 (KLRB1)		
Alternative Name:	CD161 (KLRB1 Products)		
Background:	CLEC5B, NKRP1A,NKR-P1A,HNKR-P1a,KLRB1,CD161 is a type II transmembrane glycoprotein, also known as NKR-P1A, that is expressed as a 40-44 kD homodimer. It is a member of the C-		
	type lectin superfamily. CD161 is expressed on a majority of NK cells, NKT cells, and subsets of		
	peripheral T cells and CD3+ thymocytes. It has been reported that Th17 cells are a		
	subpopulation of CD4+CD161+CCR6+ cells. While the biological function of CD161 is not clear,		
	it has been suggested to serve either as a stimulatory receptor or to inhibit NK cell-mediated		

Target Details

Storage Comment:

rarget betails		
	cytotoxicity and cytokine production. LLT-1 (lectin-like transcript-1, also named as osteoclast inhibitory lectin or CLEC2D) is the ligand of CD161.	
Gene ID:	3820	
UniProt:	Q12918	
Application Details		
Application Notes:	The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in	
	100 μL staining volume or per 100 μL of whole blood). Check your vial before the experiment.	
	Since applications vary, the appropriate dilutions must be determined for individual use.	
Comment:	Matching Isotype Control: APC Mouse IgG1, к Isotype Control (ABIN7469762)	
Restrictions:	For Research Use only	
Handling		
Buffer:	PBS with 0.05 % Proclin300, 1 % BSA	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C	

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.