

Datasheet for ABIN7456684
anti-CD40 Ligand antibody (Biotin)



[Go to Product page](#)

Overview

Quantity:	25 µg
Target:	CD40 Ligand (CD40LG)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD40 Ligand antibody is conjugated to Biotin
Application:	Flow Cytometry (FACS)

Product Details

Clone:	5C8
Isotype:	IgG2a, kappa

Target Details

Target:	CD40 Ligand (CD40LG)
Alternative Name:	CD154 (CD40LG Products)
Background:	CD40L, gp39, TRAP, T-BAM, TNFSF5, CD154 (CD40 ligand) is also known as CD40L, gp39, TRAP and T-BAM. CD40 ligand is a 32-39 kD type II transmembrane glycoprotein. It is a member of the TNF superfamily and is expressed on activated T cells. It has been reported to be important for B cell costimulation following binding of its receptor, CD40. Additionally, binding of CD40L to CD40 on B cells promotes the secretion of immunoglobulin and Ig isotype switching. CD40L is also involved in the regulation of cytokine production by T cells.

Target Details

Gene ID:	959
UniProt:	P29965
Pathways:	NF-kappaB Signaling , Production of Molecular Mediator of Immune Response , Cancer Immune Checkpoints

Application Details

Application Notes:	For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Comment:	Matching Isotype Control: Biotin Mouse Rat IgG2a, κ Isotype Control (ABIN7469786)
Restrictions:	For Research Use only

Handling

Concentration:	0.5 mg/mL
Buffer:	PBS with 0.05 % Proclin300, 1 % BSA
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2~8°C. Do not freeze.