

## Datasheet for ABIN7456734 anti-SELPLG antibody (FITC)



Overview	
Quantity:	25 µg
Target:	SELPLG
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This SELPLG antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)
Product Details	
Clone:	4RA10
Isotype:	IgG1, kappa
Target Details	
Target:	SELPLG
Alternative Name:	CD162 (SELPLG Products)
Background:	<ul> <li>P-selectin glycoprotein ligand 1,Selplg,PSGL-1,Selectin P ligand,CD162,The 4RA10 antibody</li> <li>reacts with the N-terminal functional peptide of CD162 (P-selectin glycoprotein ligand-1, PSGL-1), encoded by the Selpl gene. PSGL-1 is expressed on the cell surface as a homodimer of</li> <li>approximately 230 kDa. In the mouse, Selpl mRNA is detected in most tissues, with high levels</li> <li>found in hematopoietic cells, brain, and adipose tissue. Flow cytometric analyses have revealed</li> <li>CD162 expression on bone marrow-derived mast and dendritic cells, splenic leukocytes,</li> </ul>

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7456734 | 07/24/2024 | Copyright antibodies-online. All rights reserved.

UniProt:	Q62170
Gene ID:	20345
	tissues, and responses to vascular injury. It is a sialomucin that must be specifically sialylated, fucosylated, and sulfated to bind P-selectin. There is also evidence that other ligands for PSGL- 1 and CD62P may exist. 4RA10 mAb is reported to block the binding of mouse leukocytes to CD62P and CD62L.
	platelets, peripheral blood neutrophils, and neutrophil and T-cell lines. PSGL-1 is a ligand for P- selectin (CD62P) and is involved in leukocyte rolling, the migration of leukocytes into inflamed

## Application Details

Application Notes:	Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu$ g/10 <sup>6</sup> cells in 100 $\mu$ L volume].
Comment:	Matching Isotype Control: FITC Rat IgG1, к Isotype Control (ABIN7469821)
Restrictions:	For Research Use only
Handling	
Concentration:	0.5 mg/mL
Buffer:	PBS with 0.05 % Proclin300. 1 % BSA

Butter:	PBS WITH U.U5 % Proclin300, 1 % BSA
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.