

Datasheet for ABIN7456737 anti-ITGA4 antibody (FITC)



Overview

Quantity:	20 tests
Target:	ITGA4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ITGA4 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Clone:	BU49
Isotype:	IgG1, kappa

Target Details

Target:

ITGA4

rarget.	110/4
Alternative Name:	CD49d (ITGA4 Products)
Background:	Integrin alpha-4,ITGA4,CD49 antigen-like family member D,Integrin alpha-IV,VLA-4 subunit
	alpha,CD49d,CD49d is a 150 kD α integrin chain known as $\alpha4$ integrin or VLA-4 α chain. It forms
	a heterodimer with either integrin $\beta1$ ($\alpha4\beta1$, VLA-4) or $\beta7$ ($\alpha4\beta7$). CD49d is expressed broadly
	on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells,
	NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells,
	platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin. $\alpha4\beta7$ is the receptor

Target Details	
	for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.
Gene ID:	3676
UniProt:	P13612
Pathways:	Integrin Complex
Application Details	
Application Notes:	The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
Comment:	Matching Isotype Control: FITC Mouse IgG1, κ Isotype Control (ABIN7469760)
Restrictions:	For Research Use only
Handling	
Buffer:	PBS with 0.05 % Proclin300. 1 % BSA

Buffer:	PBS with 0.05 % Proclin300, 1 % BSA
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.