

Datasheet for ABIN7456855 anti-SLAMF1 antibody (FITC)



Go to Product page

	er		

Quantity:	50 tests
Target:	SLAMF1
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This SLAMF1 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

Product Details

Clone:	TC15-12F12-2
Isotype:	IgG2a, lambda

Target Details

Target:	SLAMF1
Alternative Name:	CD150 (SLAMF1 Products)
Background:	Slam, Slamf1, CD 150, Signaling lymphocytic activation molecule, CD150 is a 75-95 kD member of
	the immunoglobulin superfamily, also known as SLAM (signaling lymphocyte activation
	molecule) or IPO-3. CD150, a single chain type I transmembrane molecule, is expressed on
	thymocytes, T cell subsets, B cells, dendritic cells, and endothelial cells. The expression is
	upregulated upon activation. CD150 expression has been shown to be maintained on Th1 but
	not Th2 clones. T regulatory cells express a relatively high level of CD150. Antibodies against

Target Details	
	CD150 have been shown to augment IFN- γ production by Th1 cells, especially when costimulated through the TCR. CD150 associates with the src homology 2-domain-containing protein tyrosine phosphatase SHP-2, and this association is thought to be involved in signal transduction. In combination with CD48, CD150 is a useful marker for hematopoietic stem cell studies.
Gene ID:	27218
UniProt:	Q9QUM4
Application Details	
Application Notes:	The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
Comment:	Matching Isotype Control: FITC Rat IgG2a, λ Isotype Control (ABIN7469698)
Restrictions:	For Research Use only
Handling	
Buffer:	PBS with 0.05 % Proclin300. 1 % BSA

Buffer:	PBS with 0.05 % Proclin300, 1 % BSA	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.	