

Datasheet for ABIN7457295
anti-ITGA4 antibody (PE-Cy5)



[Go to Product page](#)

Overview

Quantity:	20 tests
Target:	ITGA4
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ITGA4 antibody is conjugated to PE-Cy5
Application:	Flow Cytometry (FACS)

Product Details

Clone:	9F10
Isotype:	IgG1, kappa

Target Details

Target:	ITGA4
Alternative Name:	CD49d (ITGA4 Products)
Background:	Integrin alpha-4,CD49 antigen-like family member D,Integrin alpha-IV,VLA-4 subunit alpha,CD49d,ITGA4,CD49d is a 150 kD α integrin chain known as $\alpha 4$ integrin or VLA-4 α chain. It forms a heterodimer with either integrin $\beta 1$ ($\alpha 4\beta 1$, VLA-4) or $\beta 7$ ($\alpha 4\beta 7$). CD49d is expressed broadly on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin. $\alpha 4\beta 7$ is

Target Details

the receptor for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.

Gene ID: 3676

UniProt: [P13612](#)

Pathways: [Integrin Complex](#)

Application Details

Application Notes: The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Comment: Matching Isotype Control: PE/Cyanine5 Mouse IgG1, κ Isotype Control

Restrictions: For Research Use only

Handling

Buffer: PBS with 0.05 % Proclin300, 1 % BSA

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.