

Datasheet for ABIN7457749
anti-CD57 antibody (PerCP-Cy5.5)



[Go to Product page](#)

Overview

Quantity:	20 tests
Target:	CD57 (B3GAT1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD57 antibody is conjugated to PerCP-Cy5.5
Application:	Flow Cytometry (FACS)

Product Details

Clone:	HNK-1
Isotype:	IgM, kappa

Target Details

Target:	CD57 (B3GAT1)
Alternative Name:	CD57 (B3GAT1 Products)
Background:	HNK-1, NK-1, Leu-7,CD57, also known as HNK-1, NK-1, and Leu-7 is a 100-115 kD oligosaccharide antigenic determinant expressed on a variety of proteins, lipids, and chondroitin sulfate proteoglycans. CD57 is expressed on a subset of peripheral blood lymphocytes, including NK cells and CD8+ T cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red blood cells, granulocytes, monocytes, or platelets. While the function of CD57 is unknown, binding to L-selectin, P-selectin, and a fragment of laminin

Target Details

	suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).
Gene ID:	27087
Pathways:	Glycosaminoglycan Metabolic Process

Application Details

Application Notes:	The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
Comment:	Matching Isotype Control: PerCP/Cyanine5.5 Mouse IgM, κ Isotype Control (ABIN7469749)
Restrictions:	For Research Use only

Handling

Buffer:	PBS with 0.05 % Proclin300, 1 % BSA
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.