

Datasheet for ABIN7457804 anti-ITGA4 antibody (PerCP-Cy5.5)



\sim				
()\	/e	r\ /	\cap	٨
()	<i>v</i> \leftarrow	ı vı	\Box	٧V

Quantity:20 testsTarget:ITGA4Reactivity:HumanHost:MouseClonality:MonoclonalConjugate:This ITGA4 antibody is conjugated to PerCP-Cy5.5Application:Flow Cytometry (FACS)Product DetailsClone:9F10	Overview	
Reactivity: Human Host: Mouse Clonality: Monoclonal Conjugate: This ITGA4 antibody is conjugated to PerCP-Cy5.5 Application: Flow Cytometry (FACS) Product Details	Quantity:	20 tests
Host: Mouse Clonality: Monoclonal Conjugate: This ITGA4 antibody is conjugated to PerCP-Cy5.5 Application: Flow Cytometry (FACS) Product Details	Target:	ITGA4
Clonality: Monoclonal Conjugate: This ITGA4 antibody is conjugated to PerCP-Cy5.5 Application: Flow Cytometry (FACS) Product Details	Reactivity:	Human
Conjugate: This ITGA4 antibody is conjugated to PerCP-Cy5.5 Application: Flow Cytometry (FACS) Product Details	Host:	Mouse
Application: Flow Cytometry (FACS) Product Details	Clonality:	Monoclonal
Product Details	Conjugate:	This ITGA4 antibody is conjugated to PerCP-Cy5.5
	Application:	Flow Cytometry (FACS)
Clone: 9F10	Product Details	
	Clone:	9F10

Clone:	9F10
Isotype:	IgG1, kappa

Target Details

Target:	ITGA4
Alternative Name:	CD49d (ITGA4 Products)
Background:	Integrin alpha-4,CD49 antigen-like family member D,Integrin alpha-IV,VLA-4 subunit
	alpha,CD49d,ITGA4,CD49d is a 150 kD α integrin chain known as α 4 integrin or VLA-4 α chain. It
	forms a heterodimer with either integrin $\beta1$ ($\alpha4\beta1$, VLA-4) or $\beta7$ ($\alpha4\beta7$). CD49d is expressed
	broadly on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils,
	mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red
	blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin. $\alpha4\beta7$ is

Target Details	
	the receptor for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.
Gene ID:	3676
UniProt:	P13612
Pathways:	Integrin Complex
Application Details	
Application Notes:	The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
Comment:	Matching Isotype Control: PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control (ABIN7469766)
Restrictions:	For Research Use only
Handling	
Buffer:	PBS with 0.05 % Proclin300, 1 % BSA

Buffer:	PBS with 0.05 % Proclin300, 1 % BSA
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.