antibodies -online.com





anti-CRTC2 antibody (pSer171) (Cy5.5)



Go to Product page

\sim	
()\/△	rview
\circ	1 410 44

Quantity:	100 μL
Target:	CRTC2
Binding Specificity:	pSer171
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRTC2 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Flow Cytometry (FACS)
Product Details	
- Toddet Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human CREB regulated transcription
	KLH conjugated synthetic phosphopeptide derived from human CREB regulated transcription coactivator 2 around the phosphorylation site of Ser171
Immunogen:	coactivator 2 around the phosphorylation site of Ser171
Immunogen: Isotype:	coactivator 2 around the phosphorylation site of Ser171
Immunogen: Isotype: Cross-Reactivity:	coactivator 2 around the phosphorylation site of Ser171 IgG Human, Mouse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	coactivator 2 around the phosphorylation site of Ser171 IgG Human, Mouse Rat,Dog,Pig
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	coactivator 2 around the phosphorylation site of Ser171 IgG Human, Mouse Rat,Dog,Pig

Target Details

901 2 0140	
Background:	Synonyms: TORC2, TORC-2, CREB-regulated transcription coactivator 2, Transducer of
	regulated cAMP response element-binding protein 2, Transducer of CREB protein 2, CRTC2
	Background: Transcriptional coactivator for CREB1 which activates transcription through both
	consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the
	SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of
	CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates
	gluconeogenesis as a component of the LKB1/AMPK/TORC2 signaling pathway. Regulates the
	expression of specific genes such as the steroidogenic gene, StAR. Potent coactivator of
	PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX
	activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).
Gene ID:	200186
UniProt:	Q53ET0
Pathways:	AMPK Signaling, Carbohydrate Homeostasis
Application Details	
Application Notes:	FCM 1:20-100
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months

Expiry Date: