

Datasheet for ABIN746185 anti-STAT2 antibody (pTyr690) (Biotin)

IgG

Human, Mouse, Rat

Purified by Protein A.

Dog,Horse

Overview

Isotype:

Specificity:

Purification:

Cross-Reactivity:

Predicted Reactivity:



Go to Product page

Quantity:	100 μL
Target:	STAT2
Binding Specificity:	pTyr690
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Stat2 around the phosphorylation site of Tyr690
	phosphorylation site of Tyrozo

This phosphorylation site is homologous to that of Tyr689 in Mouse and Tyr690 in Rat.

Target Details

rarget Details			
Target:	STAT2		
Alternative Name:	Stat2 (STAT2 Products)		
Background:	Synonyms: P113, ISGF-3, STAT113, Signal transducer and activator of transcription 2, STAT2 Background: Signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with IRF9/ISGF3G to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state.		
Gene ID:	6773		
UniProt:	P52630		
Pathways:	JAK-STAT Signaling, Hepatitis C, CXCR4-mediated Signaling Events		
Application Details			
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months