

## Datasheet for ABIN7464239

## anti-elF4EBP1 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	elF4EBP1 (EIF4EBP1)	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This eIF4EBP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)	

## **Product Details**

Immunogen:	Full length human 4E-BP1 Recombinant protein.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by antigen-affinity chromatography.	
Grade:	KO Validated	

## Target Details

Target:	elF4EBP1 (EIF4EBP1)	
Alternative Name:	eukaryotic translation initiation factor 4E binding protein 1 (EIF4EBP1 Products)	
Background:	Eukaryotic translation initiation factor 4E binding protein 1 , 4E-BP1 , 4EBP1 , BP-1 , PHAS-I,This	
	gene encodes one member of a family of translation repressor proteins. The protein directly	

	interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. [provided by RefSeq]	
Molecular Weight:	13 kDa	
Gene ID:	1978	
UniProt:	Q13541	
Pathways:	MAPK Signaling, PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, Regulation of Cell Size, BCR Signaling	
Application Details		
Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.	
Comment:	Positive Control: 293T , HeLa  Validation: KO/KD	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1.09 mg/mL	
Buffer:	1XPBS ( pH 7), 20 % Glycerol, 0.025 % ProClin 300	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid	

multiple freeze-thaw cycles.