



[Go to Product page](#)

Datasheet for ABIN7464807
anti-SGCG antibody

Overview

Quantity:	100 µL
Target:	SGCG
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SGCG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human gamma Sarcoglycan. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	SGCG
Alternative Name:	sarcoglycan gamma (SGCG Products)
Background:	Sarcoglycan gamma , 35DAG , A4 , DAGA4 , DMDA , DMDA1 , LGMD2C , LGMDR5 , MAM , SCARMD2 , SCG3 , gamma-SG,This gene encodes gamma-sarcoglycan, one of several

Target Details

sarcolemmal transmembrane glycoproteins that interact with dystrophin. The dystrophin-glycoprotein complex (DGC) spans the sarcolemma and is comprised of dystrophin, syntrophin, alpha- and beta-dystroglycans and sarcoglycans. The DGC provides a structural link between the subsarcolemmal cytoskeleton and the extracellular matrix of muscle cells. Defects in the encoded protein can lead to early onset autosomal recessive muscular dystrophy, in particular limb-girdle muscular dystrophy, type 2C (LGMD2C). [provided by RefSeq, Oct 2008]

Molecular Weight: 32 kDa

Gene ID: 6445

UniProt: [Q13326](#)

Application Details

Application Notes: WB: 1:500-1:3000. Optimal dilutions/concentrations should be determined by the researcher.
Not tested in other applications.

Comment: Positive Control: mouse muscle , rat muscle
Validation: Orthogonal

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: 1XPBS (pH 7), 20 % Glycerol, 0.01 % Thimerosal

Preservative: Thimerosal (Merthiolate)

Precaution of Use: This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.