

Datasheet for ABIN7464894
anti-CALCOCO2 antibody[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CALCOCO2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CALCOCO2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human NDP52. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	Purified by antigen-affinity chromatography.
Grade:	KO Validated

Target Details

Target:	CALCOCO2
Alternative Name:	calcium binding and coiled-coil domain 2 (CALCOCO2 Products)

Target Details

Background:	Calcium binding and coiled-coil domain 2 , NDP52, The protein encoded by this gene is a subunit of nuclear domain 10 (ND10) bodies. ND10 bodies are nuclear domains appearing immunohistochemically as ten dots per nucleus. They are believed to be associated with the nuclear matrix on the basis of their resistance to nuclease digestion and salt extraction. ND10 proteins are removed from the nucleus by herpes simplex virus-1 infection and may have a role in viral life cycles. [provided by RefSeq]
Molecular Weight:	52 kDa
Gene ID:	10241
UniProt:	Q13137

Application Details

Application Notes:	WB: 1:500-1:10000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. IP: 1:100-1:500. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: Raji , HepG2 (3 µM Thapsigargin treatment for 12 hr) , HepG2 (3 µM Thapsigargin treatment for 24 hr) , Huh-7 (3 µM Thapsigargin treatment for 12 hr) , Huh-7 (3 µM Thapsigargin treatment for 24 hr) Validation: KO/KD, Orthogonal
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1.57 mg/mL
Buffer:	1XPBS pH 7, 20 % Glycerol, 0.025 % ProClin 300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.