

Datasheet for ABIN7464924

anti-Transferrin antibody



Overview

Background:

Overview	
Quantity:	100 μL
Target:	Transferrin (TF)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Transferrin antibody is un-conjugated
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunofluorescence (IF)
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human Transferrin. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	Transferrin (TF)
Alternative Name:	transferrin (TF Products)

Transferrin, HEL-S-71p, PRO1557, PRO2086, TFQTL1, The protein encoded by this gene is a

glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been

	created as a result of an ancient gene duplication event that led to generation of homologous C
	and N-terminal domains each of which binds 1 ion of ferric iron. The function of this encoded
	protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal
	cells to all proliferating cells in the body. In addition to its function in iron transport, this protein
	may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the
	removal of certain organic matter/allergins from serum. [provided by RefSeq]
Molecular Weight:	77 kDa
iono ID:	7010

Molecular Weight:	77 kDa
Gene ID:	7018
UniProt:	P02787

Pathways: Transition Metal Ion Homeostasis

Application Details

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations
	should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: rat plasma
	Validation: Orthogonal
Restrictions:	For Research Use only

Handling

F	
Format:	Liquid
Concentration:	0.51 mg/mL
Buffer:	1XPBS pH 7, 20 % Glycerol, 0.025 % ProClin 300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage
	(1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid
	multiple freeze-thaw cycles.