## antibodies -online.com





## Datasheet for ABIN7465028

## anti-FBXO6 antibody



Overview	
Quantity:	100 μL
Target:	FBX06
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXO6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human FBX06. The
	exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	FBXO6
Alternative Name:	F-box protein 6 (FBXO6 Products)
Background:	F-box protein 6 , FBG2 , FBS2 , FBX6 , Fbx6b, This gene encodes a member of the F-box protein
	family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box
	proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs

	(SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class, and its C-terminal region is highly similar to that of rat NFB42 (neural F Box 42 kDa) which may be involved in the control of the cell cycle. [provided by RefSeq]
Molecular Weight:	34 kDa
Gene ID:	26270
UniProt:	Q9NRD1
Application Details	
Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: 293T
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.