## antibodies -online.com







## anti-Spermidine Synthase antibody



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Overview		
Quantity:	100 μL	
Target:	Spermidine Synthase (SRM)	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Spermidine Synthase antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human SRM. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	Spermidine Synthase (SRM)	
Alternative Name:	spermidine synthase (SRM Products)	
Background:	Spermidine synthase, PAPT, SPDSY, SPS1, SRML1, The polyamines putrescine, spermine, and spermidine are ubiquitous polycationic mediators of cell growth and differentiation. Spermidine	
	synthase is one of four enzymes in the polyamine-biosynthetic pathway and carries out the final	

Target Details		
	step of spermidine biosynthesis. This enzyme catalyzes the conversion of putrescine to spermidine using decarboxylated S-adenosylmethionine as the cofactor. [provided by RefSeq]	
Molecular Weight:	34 kDa	
Gene ID:	6723	
UniProt:	P19623	
Application Details		
Application Notes:	WB: 1:1000-1:10000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.	
Comment:	Positive Control: 293T	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal	

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.