## antibodies -online.com





## Datasheet for ABIN7465104

## anti-SYTL4 antibody



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Overview		
Quantity:	100 μL	
Target:	SYTL4	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SYTL4 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human SYTL4. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	SYTL4	
Alternative Name:	synaptotagmin like 4 (SYTL4 Products)	
Background:	Synaptotagmin like 4 , SLP4,This gene encodes a member of the synaptotagmin like protein	
	family. Members of this family are characterized by an N-terminal Rab27 binding domain and	
	C-terminal tandem C2 domains. The encoded protein binds specific small Rab GTPases and is	

## **Target Details**

rarget Details		
	involved in intracellular membrane trafficking. This protein binds Rab27 and may be involved in inhibiting dense core vesicle exocytosis. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq]	
Molecular Weight:	76 kDa	
Gene ID:	94121	
UniProt:	Q96C24	
Pathways:	Negative Regulation of Hormone Secretion	
Application Details		
Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined	
	by the researcher. Not tested in other applications.	
Comment:	Positive Control: 293T , A431 , HeLa	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	1XPBS (pH 7), 20 % Glycerol, 0.01 % Thimerosal	
Preservative:	Thimerosal (Merthiolate)	
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	omment: Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term stora (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	