

## Datasheet for ABIN7465264

## anti-POLD1 antibody (AA 1-140)



_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	POLD1	
Binding Specificity:	AA 1-140	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This POLD1 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	Recombinant protein encoding amino acids 1-140 of the large subunit of human DNA	
Immunogen:	Recombinant protein encoding amino acids 1-140 of the large subunit of human DNA polymerase d.	
Immunogen: Clone:		
	polymerase d.	
Clone:	polymerase d. 607	
Clone: Isotype:	polymerase d. 607 IgG1	
Clone:  Isotype:  Cross-Reactivity:	polymerase d.  607  IgG1  Human, Mouse	
Clone:  Isotype:  Cross-Reactivity:  Purification:	polymerase d.  607  IgG1  Human, Mouse	
Clone: Isotype: Cross-Reactivity: Purification: Target Details	polymerase d.  607  IgG1  Human, Mouse  Protein G purified	

## **Target Details**

rarget betails			
Background:	DNA polymerase delta 1, catalytic subunit, CDC2, CRCS10, MDPL, POLD, This gene encodes the 125-kDa catalytic subunit of DNA polymerase delta. DNA polymerase delta possesses bot polymerase and 3' to 5' exonuclease activity and plays a critical role in DNA replication and repair. Alternatively spliced transcript variants have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 6. [provided by RefSeq, Mar 2012]		
Molecular Weight:	124 kDa		
Gene ID:	5424		
UniProt:	P28340		
Pathways:	Telomere Maintenance, DNA Damage Repair, DNA Replication, Chromatin Binding, Synthesis o		
Application Details			
Application Notes:	WB: 1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.		
Comment:	Positive Control: 293T , HeLa , BCL-1		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 mg/mL		
Buffer:	1XPBS (pH 7), 20 % Glycerol, No Preservative		
Preservative:	Without preservative		
Storage:	4 °C,-20 °C		
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.		