

Datasheet for ABIN7465283 anti-DYNC1LI1 antibody (N-Term)



Go to Product page

Overview	
Quantity:	100 μL
Target:	DYNC1LI1
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DYNC1LI1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the N-terminus region of human
	DYNC1LI1. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	DYNC1LI1
Alternative Name:	dynein cytoplasmic 1 light intermediate chain 1 (DYNC1LI1 Products)
Background:	Dynein cytoplasmic 1 light intermediate chain 1 , DLC-A , DNCLI1 , LIC1,Acts as one of several

non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to		
be involved in linking dynein to cargos and to adapter proteins that regulate dynein function.		
Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and		
organelles along microtubules. May play a role in binding dynein to membranous organelles or		
chromosomes. Probably involved in the microtubule-dependent transport of pericentrin. Is		
required for progress throuh the spindle assembly checkpoint. The phosphorylated form		
appears to be involved in the selective removal of MAD1L1 and MAD1L2 but not BUB1B from		
kinetochores.		

Molecular Weight:	57 kDa
Gene ID:	51143
UniProt:	Q9Y6G9

Application Details

Application Notes:	WB: 1:500-1:10000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be
	determined by the researcher. Not tested in other applications.
Comment:	Positive Control: 293T , HepG2 , mouse brain
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.