

Datasheet for ABIN7465394

anti-WDR12 antibody



0	\ / /	~ P	~\ /	10	1 A
	\//	-1	١/	-	1/1
\sim	٧ '	\sim 1	v	-	V١

Overview			
Quantity:	100 μL		
Target:	WDR12		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This WDR12 antibody is un-conjugated		
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)		
Product Details			
Immunogen:	Recombinant protein encompassing a sequence within the center region of human WDR12. The exact sequence is proprietary.		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse		
Purification:	Purified by antigen-affinity chromatography.		
Target Details			
Target:	WDR12		
Alternative Name:	WD repeat domain 12 (WDR12 Products)		
Background:	WD repeat domain 12, YTM1, This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric conserved.		

Target Details	
	multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This protein is highly similar to the mouse WD repeat domain 12 protein at the amino acid level. The protein encoded by this gene is a component of a nucleolar protein complex that affects maturation of the large ribosomal subunit.
Molecular Weight:	48 kDa
Gene ID:	55759
UniProt:	Q9GZL7
Application Details	
Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: A431
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.53 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)

Preservative: Thimerosal (Merthiolate) Precaution of Use: This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.