



[Go to Product page](#)

Datasheet for ABIN746560  
**anti-CRTC1 antibody (pSer151) (Biotin)**

### Overview

Quantity:	100 µL
Target:	CRTC1
Binding Specificity:	pSer151
Reactivity:	Human, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRTC1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Torc1 around the phosphorylation site of Ser151
Isotype:	IgG
Specificity:	Due to the highly conserved nature of this protein, this antibody will react with Torc3 when phosphorylated at Ser162 across the listed species due to homology.
Cross-Reactivity:	Chicken, Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

---

Target: CRTC1

Alternative Name: Torc1 ([CRTC1 Products](#))

Background: Synonyms: MECT1, TORC1, TORC-1, WAMTP1, CREB-regulated transcription coactivator 1, Mucoepidermoid carcinoma translocated protein 1, Transducer of regulated cAMP response element-binding protein 1, Transducer of CREB protein 1, CRTC1, KIAA0616  
Background: Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR). In the hippocampus, involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses. May be required for dendritic growth of developing cortical neurons (By similarity). In concert with SIK1, regulates the light-induced entrainment of the circadian clock. In response to light stimulus, coactivates the CREB-mediated transcription of PER1 which plays an important role in the photic entrainment of the circadian clock.

Gene ID: 23373

UniProt: [Q6UUU9](#)

## Application Details

---

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

## Handling

---

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months