

[Go to Product page](#)

Datasheet for ABIN7465673 **anti-IL28B antibody**

Overview

| | |
|--------------|--------------------------------------|
| Quantity: | 100 µL |
| Target: | IL28B |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This IL28B antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|-------------------|--|
| Immunogen: | Carrier-protein conjugated synthetic peptide encompassing a sequence within the center region of human IL28B. The exact sequence is proprietary. |
| Clone: | GT727 |
| Isotype: | IgG2b |
| Cross-Reactivity: | Human |
| Purification: | Affinity purified by Protein G. |

Target Details

| | |
|-------------------|---|
| Target: | IL28B |
| Alternative Name: | interferon lambda 3 (IL28B Products) |
| Background: | Interferon lambda 3 , IFN-lambda-3 , IFN-lambda-4 , IL-28B , IL-28C , IL28B , IL28C,This gene |

Target Details

encodes a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 29 (IL29) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA). [provided by RefSeq, Jul 2008]

Molecular Weight: 22 kDa

Gene ID: 282617

UniProt: [Q8IZI9](#)

Application Details

Application Notes: WB: 1:1000-1:10000. Optimal dilutions/concentrations should be determined by the researcher.
Not tested in other applications.

Comment: Positive Control: GFP-human IL28-transfected 293T cells

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS, No Preservative

Preservative: Without preservative

Storage: 4 °C, -20 °C

Storage Comment: Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.