

Datasheet for ABIN7465700

anti-DAP3 antibody



_				
()	ve.	rv/	101	Λ

Overview		
Quantity:	100 μL	
Target:	DAP3	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DAP3 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human DAP3. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	DAP3	
Alternative Name:	death associated protein 3 (DAP3 Products)	
Background:	Death associated protein 3, DAP-3, MRP-S29, MRPS29, S29mt, bMRP-10, Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis	

within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75 % protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that also participates in apoptotic pathways which are initiated by tumor necrosis factor-alpha, Fas ligand, and gamma interferon. This protein potentially binds ATP/GTP and might be a functional partner of the mitoribosomal protein S27. Splice variants that differ in the 5' UTR have been found for this gene, both variants encode the same protein. Pseudogenes corresponding to this gene are found on chromosomes 1q and 2q. [provided by RefSeq]

Molecular Weight:	46 kDa	
Gene ID:	7818	
UniProt:	P51398	

Application Details

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations	
	should be determined by the researcher. Not tested in other applications.	
Comment:	Positive Control: H1299	
Restrictions:	For Research Use only	

Handling

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal	
Preservative:	Thimerosal (Merthiolate)	
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage	

(1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.