

Datasheet for ABIN7465769

anti-FBLIM1 antibody



()	ve	r\/i	۱۸/
\cup	V C	1 / 1	 v v

Overview			
Quantity:	100 μL		
Target:	FBLIM1		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This FBLIM1 antibody is un-conjugated		
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))		
Product Details			
Immunogen:	Recombinant protein encompassing a sequence within the center region of human FBLIM1.		
	The exact sequence is proprietary.		
Isotype:	IgG		
Cross-Reactivity:	Human		
Purification:	Purified by antigen-affinity chromatography.		
Target Details			
Target:	FBLIM1		
Alternative Name:	filamin binding LIM protein 1 (FBLIM1 Products)		
Background:	Filamin binding LIM protein 1 , CAL , FBLP-1 , FBLP1, This gene encodes a protein with an N-		
	terminal filamin-binding domain, a central proline-rich domain, and, multiple C-terminal LIM		
	domains. This protein localizes at cell junctions and may link cell adhesion structures to the		

Target Details	
	actin cytoskeleton. This protein may be involved in the assembly and stabilization of actin-filaments and likely plays a role in modulating cell adhesion, cell morphology and cell motility. This protein also localizes to the nucleus and may affect cardiomyocyte differentiation after binding with the CSX/NKX2-5 transcription factor. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq]
Molecular Weight:	41 kDa
Gene ID:	54751
UniProt:	Q8WUP2
Application Details	
Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: HCT116
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.8 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)

This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE

Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage

(1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

which should be handled by trained staff only.

4 °C,-20 °C

multiple freeze-thaw cycles.

Precaution of Use:

Storage Comment:

Storage: