## antibodies -online.com





Datasheet for ABIN7465862

## anti-Strumpellin antibody (C-Term)



| ( ) | 11/               | IN   | /ie    | A .    |
|-----|-------------------|------|--------|--------|
|     | / // <del> </del> | ۱ ات | /   (− | ' \/\/ |
|     |                   |      |        |        |

| Overview             |   |  |
|----------------------|---|--|
| Quantity:            | 100 μL  |  |
| Target:              | Strumpellin (WASHC5)  |  |
| Binding Specificity: | C-Term  |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This Strumpellin antibody is un-conjugated  |  |
| Application:         | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))  |  |
| Product Details      |   |  |
| Immunogen:           | Recombinant protein encompassing a sequence within the C-terminus region of human   |  |
|                      | Strumpellin. The exact sequence is proprietary.                                     |  |
| Isotype:             | IgG   |  |
| Cross-Reactivity:    | Human   |  |
| Purification:        | Purified by antigen-affinity chromatography.  |  |
| Target Details       |   |  |
| Target:              | Strumpellin (WASHC5)  |  |
| Alternative Name:    | WASH complex subunit 5 (WASHC5 Products)  |  |
| Background:          | WASH complex subunit 5 , KIAA0196 , RTSC , RTSC1 , SPG8,This gene encodes a 134 kDa |  |

| protein named strumpellin that is predicted to have multiple transmembrane domains and a       |  |  |
|--|--|--|
| spectrin-repeat-containing domain. This ubiquitously expressed gene has its highest expression |  |  |
| in skeletal muscle. The protein is named for Strumpell disease, a form of hereditary spastic   |  |  |
| paraplegia (HSP). Spastic paraplegias are a diverse group of disorders in which the autosomal  |  |  |
| dominant forms are characterized by progressive, lower extremity spasticity caused by axonal   |  |  |
| degeneration in the terminal portions of the longest descending and ascending corticospinal    |  |  |
| tracts. More than 30 loci (SPG1-33) have been implicated in hereditary spastic paraplegia      |  |  |
| diseases. [provided by RefSeq]   |  |  |

| Molecular Weight: | 134 kDa |
|-------------------|---------|
| Gene ID:          | 9897    |
| UniProt:          | Q12768  |

## **Application Details**

| Application Notes: | WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined |
|--------------------|--|
|                    | by the researcher. Not tested in other applications.   |
| Comment:           | Positive Control: HeLa   |
| Restrictions:      | For Research Use only  |

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | 1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal   |
| Preservative:      | Thimerosal (Merthiolate)   |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:           | 4 °C,-20 °C  |
| Storage Comment:   | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |