

[Go to Product page](#)

Datasheet for ABIN7466032

anti-PGRMC1 antibody (C-Term)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | PGRMC1 |
| Binding Specificity: | C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PGRMC1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunoprecipitation (IP) |

Product Details

| | |
|-------------------|---|
| Immunogen: | Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human PGRMC1. The exact sequence is proprietary. |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Purified by antigen-affinity chromatography. |

Target Details

| | |
|-------------------|--|
| Target: | PGRMC1 |
| Alternative Name: | progesterone receptor membrane component 1 (PGRMC1 Products) |

Target Details

| | |
|-------------------|--|
| Background: | Progesterone receptor membrane component 1 , Dap1 , HPR6.6 , IZA , MPR, Progesterone binding protein is a putative steroid membrane receptor. The protein is expressed predominantly in the liver and kidney. [provided by RefSeq] |
| Molecular Weight: | 22 kDa |
| Gene ID: | 10857 |
| UniProt: | O00264 |

Application Details

| | |
|--------------------|---|
| Application Notes: | WB: 1:500-1:3000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications. |
| Comment: | Positive Control: HepG2 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | 0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal |
| Preservative: | Thimerosal (Merthiolate) |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |