# antibodies -online.com







# anti-PDE11A antibody



| ( ) | ve     | K\ / |    | A . |
|-----|--------|------|----|-----|
|     | $\cup$ | 1 V/ | Щ. | V۷  |
|     |        |      |    |     |

| Overview          |   |  |
|-------------------|---|--|
| Quantity:         | 100 μL  |  |
| Target:           | PDE11A  |  |
| Reactivity:       | Human   |  |
| Host:             | Rabbit  |  |
| Clonality:        | Polyclonal  |  |
| Conjugate:        | This PDE11A antibody is un-conjugated   |  |
| Application:      | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))  |  |
| Product Details   |   |  |
| Immunogen:        | Recombinant protein encompassing a sequence within the center region of human PDE11A.  The exact sequence is proprietary. |  |
| Isotype:          | IgG   |  |
| Cross-Reactivity: | Human   |  |
| Purification:     | Purified by antigen-affinity chromatography.  |  |
| Target Details    |   |  |
| Target:           | PDE11A  |  |
| Alternative Name: | phosphodiesterase 11A (PDE11A Products)   |  |
| Background:       | Phosphodiesterase 11A, PPNAD2, The 3', 5'-cyclic nucleotides cAMP and cGMP function as                                    |  |
|                   | second messengers in a wide variety of signal transduction pathways. 3',5'-cyclic nucleotide                              |  |
|                   | phosphodiesterases (PDEs) catalyze the hydrolysis of cAMP and cGMP to the corresponding 5'-                               |  |

#### **Target Details**

|                   | monophosphates and provide a mechanism to downregulate cAMP and cGMP signaling. This            |
|-------------------|---|
|                   | gene encodes a member of the PDE protein superfamily. Mutations in this gene are a cause of     |
|                   | Cushing disease and adrenocortical hyperplasia. Multiple transcript variants encoding different |
|                   | isoforms have been found for this gene. [provided by RefSeq]                                    |
| Molecular Weight: | 105 kDa   |
| Gene ID:          | 50940   |
| UniProt:          | Q9HCR9  |
| Pathways:         | cAMP Metabolic Process  |

## **Application Details**

| Application Notes: | WB: 1:1000-1:10000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be |  |
|--------------------|---|--|
|                    | determined by the researcher. Not tested in other applications.                     |  |
| Comment:           | Positive Control: HepG2   |  |
| Restrictions:      | For Research Use only   |  |

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | 0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal   |
| Preservative:      | Thimerosal (Merthiolate)   |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:           | 4 °C,-20 °C  |
| Storage Comment:   | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |