

#### Datasheet for ABIN7466048

# anti-RBP4 antibody



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μL
Target:	RBP4
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RBP4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human RBP4. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	RBP4
Alternative Name:	retinol binding protein 4 (RBP4 Products)
Background:	Retinol binding protein 4, MCOPCB10, RDCCAS, This protein belongs to the lipocalin family and is the specific carrier for retinol (vitamin A alcohol) in the blood. It delivers retinol from the liver

### **Target Details**

Molecular Weight:	stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin which prevents its loss by filtration through the kidney glomeruli. A deficiency of vitamin A blocks secretion of the binding protein posttranslationally and results in defective delivery and supply to the epidermal cells. [provided by RefSeq]
Gene ID:	5950
UniProt:	P02753
Pathways:	Regulatory RNA Pathways, Positive Regulation of Peptide Hormone Secretion, Carbohydrate  Homeostasis, Production of Molecular Mediator of Immune Response

## Application Details

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations
	should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: HepG2
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.43 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.