

Datasheet for ABIN7467170
anti-STAT3 antibody (pTyr705)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	STAT3
Binding Specificity:	pTyr705
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Immunogen:	Carrier-protein conjugated synthetic peptide surrounding phospho Tyr705 of human STAT3. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Zebrafish (Danio rerio)
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	STAT3
Alternative Name:	signal transducer and activator of transcription 3 (STAT3 Products)

Target Details

Background:	Signal transducer and activator of transcription 3 , ADMIO , ADMIO1 , APRF , HIES,The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq]
Molecular Weight:	88 kDa
Gene ID:	6774
UniProt:	P40763
Pathways:	JAK-STAT Signaling , RTK Signaling , Interferon-gamma Pathway , Neurotrophin Signaling Pathway , Dopaminergic Neurogenesis , Response to Growth Hormone Stimulus , Carbohydrate Homeostasis , Stem Cell Maintenance , Hepatitis C , Protein targeting to Nucleus , Feeding Behaviour , CXCR4-mediated Signaling Events , Signaling of Hepatocyte Growth Factor Receptor

Application Details

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Validation: Orthogonal
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS (pH 7), 1 % BSA, 20 % Glycerol, 0.025 % ProClin 300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.