

## Datasheet for ABIN7467179

## anti-Caspase 1 antibody (C-Term)



Go to Product page

Overview
----------

Quantity:	100 μL
Target:	Caspase 1 (CASP1)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 1 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the C-terminus region of human
	Recombinant protein encompassing a sequence within the C-terminus region of human Caspase 1 p10 subunit. The exact sequence is proprietary.
Immunogen:	Caspase 1 p10 subunit. The exact sequence is proprietary.
Immunogen: Isotype:	Caspase 1 p10 subunit. The exact sequence is proprietary.
Immunogen:  Isotype:  Cross-Reactivity:  Purification:	Caspase 1 p10 subunit. The exact sequence is proprietary.  IgG  Human, Mouse, Rat
Immunogen:  Isotype:  Cross-Reactivity:	Caspase 1 p10 subunit. The exact sequence is proprietary.  IgG  Human, Mouse, Rat
Immunogen:  Isotype:  Cross-Reactivity:  Purification:	Caspase 1 p10 subunit. The exact sequence is proprietary.  IgG  Human, Mouse, Rat
Immunogen:  Isotype:  Cross-Reactivity:  Purification:  Target Details	Caspase 1 p10 subunit. The exact sequence is proprietary.  IgG  Human, Mouse, Rat  Purified by antigen-affinity chromatography.

aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role
in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo
proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small,
that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically
cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes
such as inflammation, septic shock, and wound healing. This gene has been shown to induce
cell apoptosis and may function in various developmental stages. Studies of a similar gene in
mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing of this
gene results in five transcript variants encoding distinct isoforms. [provided by RefSeq]

Molecular Weight:	pro-form: 45, cleaved form: 10 kDa
Gene ID:	834
UniProt:	P29466
Pathways:	Apoptosis, Interferon-gamma Pathway, Positive Regulation of Endopeptidase Activity,
	Inflammasome

## **Application Details**

Application Notes:	IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher.
	Not tested in other applications.
Comment:	Validation: Orthogonal
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS (pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.