antibodies -online.com





anti-PEX19 antibody



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	PEX19
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PEX19 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human PEX19. The exact sequence is proprietary.
Clone:	GT533
Isotype:	lgG2b
Cross-Reactivity:	Human, Rat
Purification:	Affinity purified by Protein G.

Target Details

Target:	PEX19	
Alternative Name:	tive Name: peroxisomal biogenesis factor 19 (PEX19 Products)	

Target Details

Background:	Peroxisomal biogenesis factor 19 , D1S2223E , HK33 , PBD12A , PMP1 , PMPI , PXF ,	
	PXMP1,This gene is necessary for early peroxisomal biogenesis. It acts both as a cytosolic	
	chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Peroxins	
	(PEXs) are proteins that are essential for the assembly of functional peroxisomes. The	
	peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal	
	recessive, lethal diseases characterized by multiple defects in peroxisome function. The	
	peroxisomal biogenesis disorders are a heterogeneous group with at least 14 complementation	
	groups and with more than 1 phenotype being observed in cases falling into particular	
	complementation groups. Although the clinical features of PBD patients vary, cells from all PBD	
	patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins	
	into the organelle. Defects in this gene are a cause Zellweger syndrome (ZWS). [provided by	
	RefSeq]	
Molecular Weight:	33 kDa	
Gene ID:	5824	
UniProt:	P40855	
Application Details		
Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. FACS: 1:50-1:200. Optimal dilutions/concentrations	
	should be determined by the researcher. Not tested in other applications.	
Comment:	Positive Control: PC-12 , Rat-2 , Jurkat , Raji , K562 , THP-1 , NCI-H929	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1.17 mg/mL	
Buffer:	PBS, No Preservative	
Preservative:	Without preservative	
Storage:	4 °C,-20 °C	
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage	
	(1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid	
	multiple freeze-thaw cycles.	