

### Datasheet for ABIN7467469

# anti-RGS2 antibody



Go to Product page

Overviev	

Overview	
Quantity:	100 μL
Target:	RGS2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RGS2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP)
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human RGS2. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	RGS2
Alternative Name:	regulator of G protein signaling 2 (RGS2 Products)
Background:	Regulator of G protein signaling 2, G0S8,Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of

### **Target Details**

	the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 2 belongs to this family. The protein acts as a mediator of myeloid differentiation and may play a role in leukemogenesis. [provided by RefSeq]
Molecular Weight:	24 kDa
Gene ID:	5997
UniProt:	P41220
Pathways:	Myometrial Relaxation and Contraction, Regulation of G-Protein Coupled Receptor Protein Signaling, Brown Fat Cell Differentiation

# **Application Details**

Application Notes:	WB: 1:500-1:3000. IP: 1:100-1:500. Optimal dilutions/concentrations should be determined by
	the researcher. Not tested in other applications.
Comment:	Positive Control: U87-MG
Restrictions:	For Research Use only

# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.