antibodies -online.com







anti-CYFIP1 antibody (C-Term)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Overview		
Quantity:	100 μL	
Target:	CYFIP1	
Binding Specificity:	C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CYFIP1 antibody is un-conjugated	
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the C-terminus region of human	
	CYFIP1. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	CYFIP1	
Alternative Name:	cytoplasmic FMR1 interacting protein 1 (CYFIP1 Products)	
Background:	Cytoplasmic FMR1 interacting protein 1, P140SRA-1, SHYC, SRA-1, SRA1, Component of the	

Target Details		
	CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression. In the CYFIP1-EIF4E-FMR1 complex this subunit is an adapter between EIF4E and FMR1. Promotes the translation repression activity of FMR1 in brain probably by mediating its association with EIF4E and mRNA (By similarity). Involved in formation of membrane ruffles and lamellipodia protrusions and in axon outgrowth. Binds to F-actin but not to RNA. Regulator of epithelial morphogenesis. May act as an invasion suppressor in cancers.	
Molecular Weight:	145 kDa	
Gene ID:	23191	
UniProt:	Q7L576	
Application Details		
Application Notes:	IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal	
Preservative:	Thimerosal (Merthiolate)	

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal	
Preservative:	Thimerosal (Merthiolate)	
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	