## antibodies -online.com





## Datasheet for ABIN7467937

## anti-ARL13B antibody



| $\sim$ |    |    |    |             |   |
|--------|----|----|----|-------------|---|
| ()     | VE | ۲۱ | /1 | $\triangle$ | Λ |

| Quantity:         | 100 μL  |
|-------------------|---|
| Target:           | ARL13B  |
| Reactivity:       | Human   |
| Host:             | Rabbit  |
| Clonality:        | Polyclonal  |
| Conjugate:        | This ARL13B antibody is un-conjugated   |
| Application:      | Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC),            |
|                   | Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))                           |
| Product Details   |   |
| Immunogen:        | Recombinant protein encompassing a sequence within the center region of human ARL13B. |
|                   | The exact sequence is proprietary.  |
| Isotype:          | IgG   |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Purified by antigen-affinity chromatography.  |
| Target Details    |   |

| Target:           | ARL13B   |  |
|-------------------|--|--|
| Alternative Name: | ADP ribosylation factor like GTPase 13B (ARL13B Products)                              |  |
| Background:       | Synonyms: ADP ribosylation factor like GTPase 13B , ARL2L1 , JBTS8                     |  |
|                   | Background: This gene encodes a member of the ADP-ribosylation factor-like family. The |  |

| Target Details      |   |
|---------------------|---|
|                     | encoded protein is a small GTPase that contains both N-terminal and C-terminal guanine nucleotide-binding motifs. This protein is localized in the cilia and plays a role in cilia formation and in maintenance of cilia. Mutations in this gene are the cause of Joubert syndrome 8.  Alternate splicing results in multiple transcript variants. [provided by RefSeq] |
| Molecular Weight:   | 49 kDa  |
| Gene ID:            | 200894  |
| UniProt:            | Q3SXY8  |
| Application Details |   |
| Application Notes:  | WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.   |
| Comment:            | Positive Control: Mouse liver Validation: Orthogonal  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 mg/mL   |
| Buffer:             | 0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal  |

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | 0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal   |
| Preservative:      | Thimerosal (Merthiolate)   |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:           | 4 °C,-20 °C  |
| Storage Comment:   | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |